

Pati-Salam model in curved space-time from square root Lorentz manifold

De-Sheng Li*

(Dated: June 26, 2021)

Abstract: There is a $U(4) \times U(4)$ -bundle on four-dimensional square root Lorentz manifold. Then a Pati-Salam model in curved space-time (Lagrangian) and a gravity theory (Lagrangian) are constructed on square root Lorentz manifold based on self-parallel transportation principle. An explicit formulation of Sheaf quantization on this square root Lorentz manifold is shown. Sheaf quantization is based on superposition principle and construct a linear Sheaf space in curved space-time. The transition amplitude in path integral quantization is given which is consistent with Sheaf quantization. All particles and fields in Standard Model (SM) of particle physics and Einstein gravity are found in square root metric and the connections of bundle. The interactions between particles/fields are described by Lagrangian explicitly. There are few new physics in this model. The gravity theory is Einstein-Cartan kind with torsion. There are new particles, right handed neutrinos, dark photon, Fiona, $X^{\pm C}$ and $Y^0, Y^1, Y^2, Y_*^1, Y_*^2$.

Keywords: Lorentz manifold, Pati-Salam model, Curved space-time, Gravity theory, Yang-Mills theory, Sheaf quantization

* At Guizhou Aerospace Tianma Electromechanical S&T Co.,Ltd; also at College of Physics, Mechanical and Electrical Engineering, Jishou University, Jishou 416000, P. R. China; also at Interdisciplinary Center for Quantum Information, National University of Defense Technology, Changsha 410073, P. R. China; also at Institute of High Energy Physics, Chinese Academy of Sciences, 19B Yuquan Road, Beijing 100049, P. R. China.

I. INTRODUCTION

Four-dimensional pseudo-Riemann geometry with signature $(-, +, +, +)$, Lorentz manifold, is the geometry background of the general relativity, space-time is described by the metric, and the gravitational field is described as the curve of space-time. In general relativity, the geodesic equation describes the trajectories of free particles, and the Einstein equation determines how matter curves space-time. At the last life time of Einstein, he attempted to establish a new geometry unifying electromagnetic interaction and gravity. This idea was developed by Weil into the early idea of gauge invariance and by the Kaluza and Klein into the idea of extra dimensions.

Later, the Yang-Mills theory [1] was confirmed. Yang-Mills theory takes gauge

invariance as its basic principle and to be the theoretical framework of electromagnetic, weak and strong interaction in Standard Model (SM) of particle physics [2-7]. The Yang-Mills theory is the theoretical framework of SM and has a good correspondence with the complex structure group G fiber bundle theory [8]. General relativity can actually be rewritten in the framework of fiber bundle theory also, except that the G structure group of general relativity is real, and the corresponding fiber bundles are the tangent and cotangent bundles. Another way to build unified field theory is introducing extra dimensions to give all fields their geometric positions. And lots of attempts in extra dimension were made. Is it possible to fuse the tangent (cotangent) bundle of general relativity with the complex structure group G -bundle of Yang-Mills theory?

Inspired by the Dirac's way of finding his equation and spinors through making square root of the Klein-Gordon equation, we researched the four-dimensional square root Lorentz manifold, which similar with the papers in Clifford algebra or Clifford bundle [9-52], spin-gauge theory in Riemann-Cartan

space-time [53, 54], sedenion [55] and Einstein-Cartan theory [56-58] etc. Four-dimensional square root Lorentz manifold has extra $U(4') \times U(4)$ principal bundle than Lorentz manifold. Two Lagrangians based on four-dimension square root Lorentz manifold are constructed which describe a $U(4') \times U(4)_L \times U(4)_R$ Pati-Salam model in curved space-time and a gravity theory, respectively. In the Pati-Salam model [59], the $SU(4')$ is color group with “lepton number as the fourth color”, and the $SU(4)_L \times SU(4)_R$ is chiral flavor group. The gravity theory is Einstein kind with torsion. We realize an explicit formulation of Sheaf quantization [60-76] scheme which consistent with path integral quantization. The particles spectrum on this model is analyzed.

II. GEOMETRY AND LAGRANGIAN

The notations are introduced here. a, b, c, d represent frame indices and a, b, c, d are equal to $0, 1, 2, 3$. μ, ν, ρ, σ represent coordinates indices and μ, ν, ρ, σ are equal to $0, 1, 2, 3$. α represent group indices with α equals to $0, 1, \dots, 15$. i, j, k, l, m are equal to $1, 2, 3, 4$. C is quarks color and equals to $R, G, B(1, 2, 3)$. κ is Sheaf space index. Repeated indices are summed by default.

The pseudo-Riemann manifold is described by a metric

$$g(x) = -g_{\mu\nu}(x)dx^\mu \otimes dx^\nu, \quad (1)$$

where the metric is symmetric

$$g_{\mu\nu}(x) = g_{\nu\mu}(x), \quad (2)$$

and $\{x|x = (x^\mu) = (\vec{t}, x)\}$ is a four-dimensional topological space. Here we discuss the four-dimensional pseudo-Riemann manifold with signature $(-, +, +, +)$, Lorentz manifold. And it can be described by orthonormal frame (vierbein) formalism as

$$g^{-1}(x) = -\eta^{ab}\theta_a(x)\theta_b(x), \quad (3)$$

where

$$\eta^{ab} = \text{diag}(1, -1, -1, -1).$$

Orthonormal frames describe gravitational




FIG. 1. Picture visualization of geometry background of (a) general relativity (b) Yang-Mills theory and (c) square root metric. (a) Geometry background of general relativity is pseudo-Riemann geometry, which is a smooth curved manifold. (b) Geometry background of Yang-Mills theory is complex G-bundle with flat base manifold, G is the gauge group of Yang-Mills theory. (c) Square root metric geometry in four-dimension has everything of pseudo-Riemann geometry and extra $U(4') \times U(4)$ -bundle.

field

$$\theta_a(x) = \theta_a^\mu(x) \frac{\partial}{\partial x^\mu}.$$

The definition of gamma matrices is

$$\gamma^a \gamma^b + \gamma^b \gamma^a = 2\eta_{ab} I_{4 \times 4}. \quad (4)$$

The Hermiticity conditions for gamma matrices are

$$\gamma^a \gamma^{b\dagger} + \gamma^{b\dagger} \gamma^a = 2I^{ab} I_{4 \times 4}, \quad (5)$$

where I^{ab} is $\text{diag}(1, 1, 1, 1)$. We define

$$l = i\gamma_{ik}^0(x)\gamma_{kj}^a(x)e_j^\dagger \otimes e_i\theta_a(x), \quad (6)$$

$$\tilde{l} = i\gamma_{ik}^a(x)\gamma_{kj}^0(x)e_j^\dagger \otimes e_i\theta_a(x), \quad (7)$$

where e_i are the orthogonal bases expanding four-dimension complex space \mathbb{C}^4 . The orthogonal bases of \mathbb{C}^4 satisfy

$$\text{tr}(e_j^\dagger \otimes e_i) = e_i e_j^\dagger = \delta_{ij}. \quad (8)$$

One simple choice of e_i is

$$e_1 = (e^{i\theta_1}, 0, 0, 0), e_2 = (0, e^{i\theta_2}, 0, 0), \quad (9)$$

$$e_3 = (0, 0, e^{i\theta_3}, 0), e_4 = (0, 0, 0, e^{i\theta_4}). \quad (10)$$

After using $\gamma^{a\dagger} = \gamma^0 \gamma^a \gamma^0$, we find that

$$g^{-1}(x) = \frac{1}{4} \text{tr}[\tilde{l}(x)l(x)]. \quad (11)$$

Then $l(x)$ and $\tilde{l}(x)$ are the square root of metric in some sense. The $\gamma_{ij}^a(x)$ can be shown as

$$\begin{aligned} \gamma_{ik}^{0'}(x)\gamma_{kj}^{a'}(x) &= \psi_{ik}^\dagger(x)\gamma_{kl}^0\gamma_{lm}^a\psi_{mj}(x) \\ &= \bar{\psi}_i(x)\gamma^a\psi_j(x), \\ \gamma_{ik}^{a'}(x)\gamma_{kj}^{0'}(x) &= \psi_{ik}^\dagger(x)\gamma_{kl}^a\gamma_{lm}^0\psi_{mj}(x) \\ &= \bar{\psi}_i(x)\gamma^{a\dagger}\psi_j(x), \end{aligned}$$

where ψ_i are the Dirac fermions field with flavor related index i equals to 1, 2, 3, 4. And

$$\bar{\psi}_i(x) = \psi_i^\dagger(x)\gamma^0, \psi(x) \in U(4)$$

is 4×4 matrix. So, the square root metrics are defined as follow

$$l(x) = i\bar{\psi}_i(x)\gamma^a\psi_j(x)e_j^\dagger \otimes e_i\theta_a(x), \quad (12)$$

$$\tilde{l}(x) = i\bar{\psi}_i(x)\gamma^{a\dagger}\psi_j(x)e_j^\dagger \otimes e_i\theta_a(x). \quad (13)$$

The square root Lorentz manifold is described by square root metric (12), (13). Direct calculations show that the definition (12) and (13) satisfy (11) and

$$l^\dagger(x) = -l(x), \quad \tilde{l}^\dagger(x) = -\tilde{l}(x).$$

The coefficients of the affine connections on coordinates, coefficients of spin connections on orthonormal frame [77] and gauge fields on the $U(4') \times U(4)$ -bundle are defined as follows

$$\nabla_\mu \partial_\nu = \Gamma_{\nu\mu}^\rho(x)\partial_\rho, \quad (14)$$

$$\nabla_\mu \theta_a(x) = \Gamma_{a\mu}^b(x)\theta_b(x), \quad (15)$$

$$\nabla_\mu(\gamma^0\gamma^a) = i[V_\mu(x)\gamma^0\gamma^a - \gamma^0\gamma^a V_\mu(x)], \quad (16)$$

$$\nabla_\mu e_i = iW_{\mu ij}(x)e_j^\dagger. \quad (17)$$

A relation between coefficients of the affine connections on coordinates and coefficients of spin connections on orthonormal frame can be easily found

$$\Gamma_{a\mu}^b(x)\theta_b^\rho(x) = \partial_\mu\theta_a^\rho(x) + \theta_a^\nu(x)\Gamma_{\nu\mu}^\rho(x).$$

If the covariant derivative is compatible with metric,

$$\nabla g(x) = 0,$$

then

$$\Gamma_{ab\mu} = -\Gamma_{ba\mu}.$$

The gauge fields on the $U(4') \times U(4)$ -bundle are Hermitian

$$V_\mu^\dagger(x) = V_\mu(x), W_{\mu ij}^*(x) = W_{\mu ji}(x).$$

The uniqueness of definition of gauge fields is originated from restriction (4), (5) and (8). The equation as follow can be derived from (16) $\nabla_\mu(\gamma^a\gamma^0) = i[\tilde{V}_\mu(x)\gamma^a\gamma^0 - \gamma^a\gamma^0\tilde{V}_\mu(x)]$, (18) where we write

$$\tilde{V}_\mu(x) = \gamma^0 V_\mu \gamma^0.$$

The gauge field $V_\mu(x)$ and $W_{\mu ij}(x)$ can be decomposed by the generators of the $U(4)$ group

$$V_\mu(x) = V_\mu^\alpha(x)\mathcal{T}^\alpha, \quad (19)$$

$$W_{\mu ij}(x) = W_{\mu ij}^\alpha(x)\mathcal{T}_{ij}^\alpha, \quad (20)$$

where α is equals to 0, 1, 2, ..., 15. There are Hermitian gauge bosons fields

$$V_\mu^{\alpha\dagger}(x) = V_\mu^\alpha(x), W_{\mu ij}^{\alpha\dagger}(x) = W_{\mu ij}^\alpha(x).$$

The \mathcal{T}^α are the generators of $U(4)$ and an explicit one can be seen in appendix. An equation is constructed which satisfying the $U(4') \times U(4)$ gauge invariant, locally Lorentz invariant and generally covariant principles

$$\text{tr} \nabla[l(x)] = 0. \quad (21)$$

This equation originated from generalized self-parallel transportation principle. Eliminating index x , the explicit formula of equation (21) is

$$\begin{aligned} &[(i\partial_\mu\bar{\psi}_i - \bar{\psi}_i\tilde{V}_\mu + W_{\mu ij}\bar{\psi}_j)\gamma^a\psi_i + \bar{\psi}_i\gamma^a(i\partial_\mu\psi_i \\ &\quad + V_\mu\psi_i - \psi_jW_{\mu ji}) \\ &\quad + i\bar{\psi}_i\gamma^b\psi_j\Gamma_{b\mu}^a]\theta_a = 0. \end{aligned}$$

We define a Lagrangian

$$\mathcal{L} = \bar{\psi}_i\gamma^a(i\partial_\mu\psi_i + V_\mu\psi_i - \psi_jW_{\mu ji})\theta_a^\mu + \frac{i}{2}\bar{\psi}_i\gamma^b\psi_j\Gamma_{b\mu}^a\theta_a^\mu. \quad (22)$$

The last term in Lagrangian (22) is Yukawa coupling term $\psi_i\phi\psi_i$ and the scalar (Higgs) field is gamma matrix valued and originated from gravitational field

$$\phi = \frac{i}{2}\gamma^b\Gamma_{b\mu}^a\theta_a^\mu. \quad (23)$$

Then, the Lagrangian (22) describes $U(4') \times U(4)$ Yang-Mills theory in curved space-time.

The Lagrangian (22) has relation with (21)

$$\text{tr} \nabla[l(x)] = \mathcal{L} - \mathcal{L}^\dagger. \quad (24)$$

If equation (21) being satisfied, the Lagrangian (22) is Hermitian

$$\mathcal{L} = \mathcal{L}^\dagger. \quad (25)$$

So, the unitary principle of quantum field theory (25) consistent with generalized self-parallel transportation principle (21). The equations of motion for the Lagrangian (22) are

$$\gamma^a (i\partial_\mu \psi_i + V_\mu \psi_i - \psi_j W_{\mu ji}) \theta_a^\mu + \frac{i}{2} \gamma^b \psi_i \Gamma_{b\mu}^a \theta_a^\mu = 0, \quad (26)$$

and this equation's conjugate transpose. We point out that (21) is density matrix version of (26). The effective equation of motion of (26) has signature (1, -1, -1, -1). For example, the massless Dirac equation in curved space-time of this model is

$$i\gamma^a \theta_a^\mu \partial_\mu \psi_i = 0. \quad (27)$$

The square of equation (27) is massless Klein-Gordon equation in curved space-time

$$\eta^{ab} \theta_a^\mu \theta_b^\nu \partial_\mu \partial_\nu \psi_i = 0. \quad (28)$$

The signature of equation (28) is (1, -1, -1, -1) and consistent with special relativity. Then, a Lagrangian (22) which describes the $U(4') \times U(4)$ Yang-Mills theory in curved space-time is all other fields (except Higgs field) dynamical background which satisfies the characteristic of the gravitational field in our real world.

Lagrangian (22) is $U(4') \times U(4)$ gauge invariant, locally Lorentz invariant and generally covariant. So, Lagrangian (22) is demanded invariant under the transformations

$$\begin{aligned} \psi_i' &= \tilde{U} \psi_j U_{ji}, \\ \gamma^{a'} &= \tilde{U} \gamma^b \tilde{U}^\dagger \Lambda_b^a, \\ \theta_a'^\mu &= \Lambda_b^a \theta_b^\mu, \end{aligned} \quad (29)$$

Then, the transformation rules have to be derived as follows

$$V_\mu' = \tilde{U} V_\mu \tilde{U}^\dagger - (\partial_\mu \tilde{U}) \tilde{U}^\dagger, \quad (30)$$

$$V_\mu' = U_{ki}^* W_\mu U_{lj} + U_{ki}^* (\partial_\mu U_{kj}), \quad (31)$$

$$\Gamma_{a\mu}^{b'} = \Lambda_a^c \Gamma_{c\mu}^d \Lambda_d^b - \Lambda_a^c \partial_\mu \Lambda_c^b, \quad (32)$$

where the transformation matrices satisfy

$$\tilde{U} \tilde{U}^\dagger = I, U_{ji} U_{jk} = \delta_{ik}, \Lambda_a^b \Lambda_b^c = \delta_a^c.$$

Then the transformation elements

$$\tilde{U} \in U(4'),$$

$$(U_{ij}) \in U(4),$$

$$\Lambda_b^a \in O(1,3),$$

and $U(4')$ is color group, $U(4)$ is flavor group, $O(1,3)$ is locally Lorentz group. Then we complete the proof of the Lagrangian (22) is $U(4') \times U(4)$ gauge invariant, locally Lorentz invariant and generally covariant.

The gauge field strength tensors and curvature tensor are defined as follows

$$\begin{aligned} H_{\mu\nu} &= \partial_\mu V_\nu - \partial_\nu V_\mu - iV_\mu V_\nu + iV_\nu V_\mu, \\ F_{\mu\nu ij} &= \partial_\mu W_{\nu ij} - \partial_\nu W_{\mu ij} - iW_{\mu ik} W_{\nu kj} \\ &\quad + iW_{\nu ik} W_{\mu kj}, \\ R_{b\mu\nu}^a &= \partial_\mu \Gamma_{b\nu}^a - \partial_\nu \Gamma_{b\mu}^a + \Gamma_{b\nu}^c \Gamma_{c\mu}^a \\ &\quad - \Gamma_{b\mu}^c \Gamma_{c\nu}^a. \end{aligned}$$

If the covariant derivative is compatible with metric, the curvature tensor satisfies

$$R_{ab\mu\nu} = -R_{ba\mu\nu}.$$

The gauge fields strengths on the $U(4') \times U(4)$ -bundle match the Hermiticity conditions

$$H_{\mu\nu}^\dagger = H_{\mu\nu}, F_{\mu\nu ij}^* = F_{\mu\nu ji}.$$

The gauge field strength can be decomposed by the $U(4)$ generators

$$H_{\mu\nu} = H_{\mu\nu}^\alpha \mathcal{T}_{ij}^\alpha, F_{\mu\nu ij} = F_{\mu\nu ij}^\alpha \mathcal{T}_{ij}^\alpha.$$

After the torsion being defined

$$T_{\nu\rho}^a = 2\Gamma_{[\nu\rho]}^a, \quad (33)$$

we have the Ricci identity and Bianchi identity [78] on this geometry structure as follows

$$\partial_{[\mu} H_{\nu\rho]} = H_{[\mu\nu} V_{\rho]} - V_{[\mu} H_{\nu\rho]}, \quad (34)$$

$$\partial_{[\mu} F_{\nu\rho] ij} = F_{[\mu\nu] ik} W_{\rho] kj} - W_{[\mu ik} F_{\nu\rho] kj}, \quad (35)$$

$$T_{\sigma[\rho}^a T_{\mu\nu]} = R_{[\rho\mu\nu]}^a + \nabla_{[\rho} T_{a\mu\nu]}, \quad (36)$$

$$\nabla_{[\rho} R_{b|\mu\nu]}^a = R_{b\sigma[\rho}^a T_{\mu\nu]\sigma}. \quad (37)$$

There is Yang-Mills Lagrangian for gauge bosons in this model

$$\mathcal{L}_Y = \frac{-1}{2} \text{tr}(H^{\mu\nu} H_{\mu\nu}) - \frac{-\zeta}{2} F_{ij}^{\mu\nu} F_{\mu\nu ji}, \quad (38)$$

where $\zeta \in R$ is constant.

For the gravity, the Einstein-Hilbert action in Lorentz manifold be showed as follow

$$S = \int \omega R, \quad (39)$$

where R is the Ricci scalar curvature in Lorentz manifold, ω is volume form

$$\omega = \sqrt{-g_\nu} dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3$$

and

$$g_v = \det(g_{\mu\nu}(x)).$$

In this geometry framework, the equations can be derived as follows

$$\nabla_{[\mu}\nabla_{\nu]}l = \frac{-1}{2}(\psi_i\gamma^a\psi_k F_{\mu\nu kj} - F_{\mu\nu ki}^*\bar{\psi}_k\gamma^a\psi_j + \bar{\psi}_i\tilde{H}_{\mu\nu}^{\text{Sheaf}}\gamma^a\psi_j - \bar{\psi}_i\gamma^a H_{\mu\nu}\psi_j + \frac{i}{2}\bar{\psi}_i\gamma^b\psi_j R_{b\mu\nu}^a)e_j^\dagger \otimes e_i\theta^a, \quad (40)$$

$$\nabla_{[\mu}\nabla_{\nu]}\tilde{l} = \frac{-1}{2}(\psi_i\gamma^{a\dagger}\psi_k F_{\mu\nu kj} - F_{\mu\nu ki}^*\bar{\psi}_k\gamma^{a\dagger}\psi_j + \bar{\psi}_i\tilde{H}_{\mu\nu}^{\text{Sheaf}}\gamma^{a\dagger}\psi_j - \bar{\psi}_i\gamma^{a\dagger} H_{\mu\nu}\psi_j + \frac{i}{2}\bar{\psi}_i\gamma^{b\dagger}\psi_j R_{b\mu\nu}^a)e_j^\dagger \otimes e_i\theta^a, \quad (41)$$

where we write

$$\tilde{H}_{\mu\nu} = \gamma^0 H_{\mu\nu} \gamma^0.$$

We define

$$\nabla^2 = \nabla_{[\mu}\nabla_{\nu]}dx^\mu \wedge dx^\nu,$$

the equation of a gravity theory is constructed

$$\text{tr}\nabla^2[\tilde{l}(x)l(x)] = 0. \quad (42)$$

This equation (42) is obviously $U(4') \times U(4)$ gauge invariant, locally Lorentz invariant and generally covariant. The explicit formula of equation (42) is

$$R\psi_i^\dagger\psi_i = i(F^{abij}\psi_j^\dagger(\gamma^a\gamma^b - \gamma^{b\dagger}\gamma^{a\dagger})\psi_i - \psi_i^\dagger H_{ab}(\gamma^a\gamma^b - \gamma^{b\dagger}\gamma^{a\dagger})\psi_i), \quad (43)$$

where we use the notations as follow,

$$\begin{aligned} \partial_\mu dx^\nu \otimes dx^\rho \partial_\sigma &= \delta_\mu^\nu \delta_\sigma^\rho, \\ dx^\mu \otimes dx^\nu \partial_\rho \partial_\sigma &= \delta_\rho^\mu \delta_\sigma^\nu, \\ F_{abij} &= F_{abij}\theta_a^\mu \theta_b^\nu, \\ H_{ab} &= H_{\mu\nu}\theta_a^\mu \theta_b^\nu. \end{aligned}$$

So, we define a $U(4') \times U(4)$ gauge invariant, locally Lorentz invariant, generally covariant Lagrangian

$$\begin{aligned} \mathcal{L}_g &= R\psi_i^\dagger\psi_i - i(F^{abij}\psi_j^\dagger(\gamma^a\gamma^b - \gamma^{b\dagger}\gamma^{a\dagger})\psi_i \\ &\quad + i\psi_i^\dagger H_{ab}(\gamma^a\gamma^b - \gamma^{b\dagger}\gamma^{a\dagger})\psi_i). \end{aligned} \quad (44)$$

The Lagrangian (44) is Hermitian

$$\mathcal{L}_g = \mathcal{L}_g^\dagger. \quad (45)$$

The $R\psi_i^\dagger\psi_i$ in Lagrangian (44) gives us the Einstein-Hilbert action. The equation (43) and the Einstein tensor can be derived from the Einstein-Hilbert action. The gravity theory in this model is Einstein-Cartan kind with torsion.

III. SHEAF QUANTIZATION AND PATH INTEGRAL QUANTIZATION

The entities $l(x)$ and $\tilde{l}(x)$ are two sections of the two bundles, respectively, where these two bundles are dual to each other. Further, the

Sheaf valued entities $\hat{l}(x)$ and $\hat{\tilde{l}}(x)$ are superposition of sections $l(x)$ and $\tilde{l}(x)$

$$\hat{l}(x) = \sum_\kappa \eta_\kappa(x)|\kappa, x\rangle\langle\kappa, x|l_\kappa(x), \quad (46)$$

$$\hat{\tilde{l}}(x) = \sum_\kappa \eta_\kappa(x)|\kappa, x\rangle\langle\kappa, x|\tilde{l}_\kappa(x), \quad (47)$$

where $\eta_\kappa(x) \in [0, 1]$ are probability of



FIG. 2. The fiber bundle structure is shown. Left: $l(x)$ is a section of the bundle. Right: A Sheaf is a collection of the sections. $\hat{l}(x)$ is Sheaf valued.

corresponding section $l_\kappa(x)$ and $\tilde{l}_\kappa(x)$, κ is Sheaf space index and evaluated in an abelian group. The density matrix corresponds to $\hat{l}(x)$ and $\hat{\tilde{l}}(x)$ is

$$\rho(x) = \sum_\kappa \eta_\kappa(x)|\kappa, x\rangle\langle\kappa, x|. \quad (48)$$

We have orthogonal bases in Sheaf space. The orthogonal bases in Sheaf space satisfy probability complete formulas

$$\langle\kappa, x|\kappa', x'\rangle = \delta(x - x')\delta(\kappa - \kappa'). \quad (49)$$

$$\text{tr}\rho(x) = \sum_\kappa \eta_\kappa(x) = 1. \quad (50)$$

In mathematics, a Sheaf is a collection of sections, the index κ of each section correspond to an abelian group element. In physics, the Sheaf spaces $Sh(x)$ and $\widetilde{Sh}(x)$ are expanded by all possible sections $l(x)$ and $\tilde{l}(x)$ of the two bundles, respectively. The $Sh(x)$ and $\widetilde{Sh}(x)$ are linear spaces, which means, for example, any two entities $\hat{l}_1(x)$ and $\hat{l}_2(x)$ in $Sh(x)$, there is an entity $\hat{l}(x)$ in $Sh(x)$ equals to the mixing of the two entities

$$\hat{l}(x) = \eta_1(x)\hat{l}_1(x) + \eta_2(x)\hat{l}_2(x),$$

$$\begin{aligned}\hat{l}_1(x), \hat{l}_2(x) &\in Sh(x); \\ \Rightarrow \hat{l}(x) &\in Sh(x),\end{aligned}\quad (51)$$

where the probability of each section

$$\eta_1(x), \eta_2(x) \in [0,1]$$

and

$$\eta_1(x) + \eta_2(x) = 1.$$

The Sheaf spaces $Sh(x)$ and $\widetilde{Sh}(x)$ are dual to each other. We call it Sheaf quantization which switching study objects from single section to all possible sections of the bundle. The equation (50) derives to equation of motion for density matrix

$$d(\text{tr} \rho(x)) = \text{tr}(d\rho(x)) = 0, \quad (52)$$

where d is exterior differential derivative.

The equations of motion for entities $\hat{l}(x)$ and $\hat{\tilde{l}}(x)$ in Sheaf quantization method are

$$\text{tr} \nabla[\hat{l}(x)] = 0, \text{tr}[\hat{\tilde{l}}(x)\hat{l}(x)] = 0. \quad (53)$$

The corresponding total Lagrangian density is

$$\hat{\mathcal{L}} = \sum_{\kappa} \mathcal{L}_{\kappa} + g\mathcal{L}_{g,\kappa} + \tilde{g}\mathcal{L}_{\tilde{g},\kappa}, \quad (54)$$

where g and \tilde{g} are Lagrangian multipliers with

$$g, \tilde{g} \in \mathbb{R}.$$

For pure state

$$\rho(x) = |\Psi(x)\rangle\langle\Psi(x)|, \quad (55)$$

$$|\Psi(x)\rangle = \sum_{\kappa} \alpha_{\kappa}(x)|\kappa\rangle, \quad (56)$$

where the $|\Psi(x)\rangle$ is the quantum state of quantum field theory. The transition amplitude can be defined through path integral formula

$$\alpha_{\kappa}(t, \vec{x}) = \int_{t' \in (t_0, t)} D\kappa(t', \vec{x}) e^{i\omega \mathcal{L}[\kappa(t, \vec{x})]} \alpha_{\kappa}(t_0, \vec{x}), \quad (57)$$

where ω is volume form.

IV. PARTICLES SPECTRUM

V_{μ}^{α} and W_{μ}^{α} (α is equals to $0, 1, \dots, 15$ here) are gauge bosons fields. The interactions related with the gauge bosons fields W_{μ}^{α} always preserves the possibility of chiral symmetry breaking in the Lagrangian (54) such that the gauge group can be decomposed to $U(4') \times U(4)_L \times U(4)_R$, where $U(4')$ is




FIG. 3. Weight diagram of $SU(4)$ adjoint representation and corresponding gauge bosons. The decomposition of $SU(4)$ adjoint representation is $\mathbf{15} = \mathbf{8} \oplus \mathbf{1} + \mathbf{3} + \mathbf{3}^*$. (a) The weight diagram of V_{μ}^{α} which contained gauge bosons are gluons, photon and $X^{\pm C}$. (b) The weight diagram of W_{μ}^{α} related gauge bosons are $Y^0, Y^1, Y^2, Y^*, Y^*_2$ and W^{\pm}, Z .

color group and $U(4)_L \times U(4)_R$ is chiral flavor group. The V_{μ}^0 is dark photon and W_{μ}^0 is Fiona particle. V_{μ}^0 and W_{μ}^0 are related with $U(1')$ and $U(1)$ gauge group in $U(4')$ and $U(4)$, respectively. The left-over gauge group is a Pati-Salam gauge group $SU(4') \times SU(4)_L \times SU(4)_R$ [59, 79]. The $SU(4')$ gauge group can be decomposed as follow $SU(4') = SU(3') \oplus U(1') + U_{X^+} + U_{X^-}$. (58) The $SU(3')$ is the gauge group of quantum chromodynamics (QCD) and the corresponding gauge bosons V_{μ}^{α} (α is equals to $1, 2, \dots, 8$ here) are gluons. The $U(1')$ in equation (58) is electro-magnetic interaction gauge group and corresponding gauge boson V_{μ}^{15} is photon γ . The $X^{\pm C}$ particles transport semi-leptonic processes and

$$X^{\pm C} = V_{\mu}^{8+C} \pm iV_{\mu}^{9+C}. \quad (59)$$

The electric charges of each X^{+C} and X^{-C} are $1/3$ and $-1/3$, respectively. The chiral gauge group $SU(4)_{L,R}$ can be decomposed as

$$SU(4)_{L,R} = SU(3)_Y \oplus U(1)_Z + U_{W^+} + U_{W^-}, \quad (60)$$

and related gauge bosons V_{μ}^{α} (with α equals to $1, 2, \dots, 15$) contain weak bosons W^{\pm} and Z

$$\begin{aligned}W_{\mu}^{\pm} &= W_{\mu}^9 \pm iW_{\mu}^{10} = W_{\mu}^{11} \pm iW_{\mu}^{12} \\ &= W_{\mu}^{13} \pm iW_{\mu}^{14},\end{aligned}\quad (61)$$

$$Z_{\mu} = W_{\mu}^{15}. \quad (62)$$




FIG. 4. Weight diagram of $SU(4)$ fundamental representation $\mathbf{4} \otimes \mathbf{6}$. Representation $\mathbf{4} = \mathbf{3} + \mathbf{1}$, $\mathbf{3}$ is 3 kinds of color, red, green and blue, $\mathbf{1}$ is the lepton number. Representation $\mathbf{6}$ gives us 6 flavors of quarks and leptons. 6 flavors have 3 generations, I, II and III, each generation has 2 kinds of quarks or leptons.

The left over gauge bosons are $Y^0, Y^1, Y^2, Y_*^1, Y_*^2$ with 0 electric charge. The gauge bosons Y^1, Y^2, Y_*^1, Y_*^2 transport non-SM flavor changing neutral currents (FCNCs) and

$$\begin{pmatrix} (2-\eta)Y_\mu^0 & Y_\mu^1 & Y_\mu^2 \\ Y_{*\mu}^1 & \eta Y_\mu^0 & Y_\mu^1 \\ Y_{*\mu}^2 & Y_{*\mu}^1 & -2Y_\mu^0 \end{pmatrix} = 2 \sum_{\alpha=1}^8 W_\mu^\alpha \mathcal{T}^\alpha. \quad (63)$$

The $X^{\pm C}$ and Y^1, Y^2, Y_*^1, Y_*^2 must be superheavy from the restrictions of experimental data.

The fermionic fields ψ_i transfer as the $U(4') \times U(4)$ fundamental representation according to (29). So, fermions are filled into the $SU(4)$ fundamental representation $\mathbf{4} \otimes \mathbf{6}$ naturally. The fundamental representation $\mathbf{4}$ corresponds to 3 colors and 1 lepton and leads us reobtain “Lepton number as the fourth color” [59]. The fundamental representation $\mathbf{6}$ corresponds to 6 flavors of quarks and leptons. The weight diagram coordinates in Chevalley basis of representation $\mathbf{6}$ (see in TABLE. I) have good correspondence with the quark quantum number [80]. The antifermions be filled into the representation $\mathbf{6} \otimes \mathbf{4}$ similarly. The weight diagram of fermions is shown in FIG. 4. Both left-handed and right-handed fermions for all quarks and leptons are existed.

Especially, the existence of right-handed neutrinos is predicted. This is compatible with experimental results [81] and the well know Seesaw mechanism [82-86]. The $2I_z$ being used in Table I are not no reasons because we have the Gell-Mann-Nishijima formula [80]

$$Q = I_z + \frac{\mathbb{B}+S+C+B+T}{2}. \quad (64)$$

An explicit fermions representation [79] in this model might be

$$\psi_i = \begin{pmatrix} \sqrt{2}u_R & \sqrt{2}c_R & \sqrt{2}t_R & d'_R \\ \sqrt{2}u_G & \sqrt{2}c_G & \sqrt{2}t_G & d'_G \\ \sqrt{2}u_B & \sqrt{2}c_B & \sqrt{2}t_B & d'_B \\ e & \mu & \tau & \nu' \end{pmatrix}, \quad (65)$$

where u, c, t and d' are quarks fields, e, μ, τ and ν' are electron, mu, tau and neutrinos fields. The corresponding fermions electric charges of (65) are

$$Q = \begin{pmatrix} \frac{2}{3} & \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \\ \frac{2}{3} & \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \\ \frac{2}{3} & \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \\ \frac{2}{3} & \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \\ 1 & 1 & 1 & 0 \end{pmatrix}. \quad (66)$$

The quarks states like $|d\rangle, |s\rangle, |b\rangle$ and neutrinos states $|\nu_e\rangle, |\nu_\mu\rangle, |\nu_\tau\rangle$ are eigen states of the Lagrangian [79].

	$2I_z$	S+C	B+T	H1	H2	H3
u	1	0	0	1	-1	1
d	-1	0	0	-1	1	-1
c	0	1	0	1	0	-1
s	0	-1	0	-1	0	1
t	0	0	1	0	1	0
b	0	0	-1	0	-1	0

TABLE I. Corresponding relations between quarks quantum number and weight diagram coordinates in Chevalley basis H1, H2 and H3 of $SU(4)$ representation $\mathbf{6}$ are shown.

V. CONCLUSION AND DISCUSSION

A Pati-Salam model and a gravity theory from square root Lorentz manifold are derived. A Sheaf quantization scheme which consistent with path integral quantization is shown. The particles spectrum in this model is discussed. Some possible new physics on this model are listed as follows. The gravity theory has torsion. There are exotic gauge bosons such as dark photon, Fiona, $X^{\pm C}$ and $Y^0, Y^1, Y^2, Y_*^1, Y_*^2$. The $X^{\pm C}$ transports semi-leptonic processes, the Y^1, Y^2, Y_*^1, Y_*^2 transport non-SM FCNCs. The right-handed neutrinos are existed. The Higgs field is gamma matrix valued.

ACKNOWLEDGMENTS

We thank Professor Chao-Guang Huang for long time discussions and powerful help. Without Professor Huang's help, this work is impossible. We thank Jun-Bao Wu, Zhao Li, Cai-Dian Lv, Ming Zhong, Yong-Chang Huang, Qi-Shu Yan, Yu Tian, De-Shan Yang, Yang-Hao Ou for valuable discussions. We thank Jean Thierry-Mieg for critical comments. We thank Yu Lu for computer drawing.

Appendix: Generators of U (4)

$$\begin{aligned}
 \mathcal{T}^1 &= \frac{1}{2} \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \mathcal{T}^2 = \frac{1}{2} \begin{pmatrix} 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \mathcal{T}^3 = \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\
 \mathcal{T}^4 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \mathcal{T}^5 = \frac{1}{2} \begin{pmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \mathcal{T}^6 = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\
 \mathcal{T}^7 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \mathcal{T}^8 = \frac{\sqrt{3}}{6} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \mathcal{T}^9 = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \\
 \mathcal{T}^{10} &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & -i \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \end{pmatrix}, \mathcal{T}^{11} = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \mathcal{T}^{12} = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -i \\ 0 & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{pmatrix}, \\
 \mathcal{T}^{13} &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \mathcal{T}^{15} = \frac{\sqrt{6}}{12} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -3 \end{pmatrix}, \mathcal{T}^0 = \frac{1}{2\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.
 \end{aligned}$$

Reference

- [1] C. N. Yang and R. L. Mills, "Conservation of Isotopic Spin and Isotopic Gauge Invariance," Physical Review 96, 191-195 (1954).
- [2] S. Glashow, "Partial Symmetries of Weak Interactions," Nucl.Phys. 22, 579-588 (1961).
- [3] S. Weinberg, "A Model of Leptons," Phys.Rev.Lett. 19, 1264-1266 (1967).
- [4] A. Salam, "Weak and Electromagnetic Interactions," Conf.Proc. C680519, 367-377 (1968).
- [5] D. J. Gross and F. Wilczek, "Ultraviolet Behavior of Nonabelian Gauge Theories," Phys.Rev.Lett. 30, 1343-1346 (1973).
- [6] H. D. Politzer, "Reliable Perturbative Results for Strong Interactions?" Phys.Rev.Lett. 30, 1346-1349 (1973).
- [7] P. Langacker, "Grand Unified Theories and Proton Decay," Phys.Rept. 72, 185 (1981).
- [8] B.-Y. Hou and B.-Y. Hou, Differential geometry for physicists (1999).
- [9] J. Dowker and R. Banach, "Quantum Field Theory on Clifford-klein Space-times. The

- Effective Lagrangian and Vacuum Stress Energy Tensor,” *J. Phys. A* 11, 2255 (1978).
- [10] R. Casalbuoni and R. Gatto, “Unified Theories for Quarks and Leptons Based on Clifford Algebras,” *Phys. Lett. B* 90, 81-86 (1980).
- [11] W. Rodrigues and Q. de Souza, “The Clifford bundle and the nature of the gravitational field,” *Found. Phys.* 23, 1465-1490 (1993).
- [12] J. Smith, Frank D. Tony, “SU(3) x SU(2) x U(1), Higgs, and gravity from spin (0,8) Clifford algebra Cl(0,8),” (1994), arXiv:hep-th/9402003.
- [13] H. White and R. Lloyd, “Gauge field theories on Clifford algebras,” *Lett. Math. Phys.* 41, 349-370 (1997).
- [14] J. Keller, “Twistors and Clifford Algebras,” *Fundam. Theor. Phys.* 94, 161-173 (1998).
- [15] J. Vargas and D. Torr, “Marriage of Clifford algebra and Finsler geometry: A lineage for unification?” *Int. J. Theor. Phys.* 40, 275-298 (2001).
- [16] M. Pavsic, “Clifford space as the arena for physics,” *Found. Phys.* 33, 1277-1306 (2003), arXiv:gr-qc/0211085.
- [17] W. Krolkowski, “Clifford algebra implying three fermion generations revisited,” *Acta Phys. Polon. B* 33, 2559-2570 (2002), arXiv:hep-ph/0203107.
- [18] R. A. Mosna and J. Rodrigues, Waldyr A., “The bundles of algebraic and dirac-hestenes spinor fields,” *J. Math. Phys.* 45, 2945-2966 (2004), arXiv:math-ph/0212033 [math-ph].
- [19] C. Castro and M. Pavsic, “Clifford algebra of space-time and the conformal group,” *Int. J. Theor. Phys.* 42, 1693-1705 (2003), arXiv:hep-th/0203194.
- [20] M. El Naschie, “Quantum gravity, Clifford algebras, fuzzy set theory and the fundamental constants of nature,” *Chaos Solitons Fractals* 20, 437-450 (2004).
- [21] M. Pavsic, “Clifford space as a generalization of spacetime: Prospects for unification in physics,” in 4th Vigier: The Search for Unity in Physics: A Symposium in Celebration of the 83rd Birthday of Mathematical Physicist Jean-Pierre Vigier (2004) arXiv:hep-th/0411053.
- [22] M. Pavsic, “Kaluza-Klein theory without extra dimensions: Curved Clifford space,” *Phys. Lett. B* 614, 85-95 (2005), arXiv:hep-th/0412255.
- [23] M. Pavsic, “Spin gauge theory of gravity in Clifford space: A Realization of Kaluza-Klein theory in 4-dimensional spacetime,” *Int. J. Mod. Phys. A* 21, 5905-5956 (2006), arXiv:gr-qc/0507053.
- [24] S. Vacaru, P. Stavrinou, E. Gaburov, and D. Gonta, Clifford and Riemann-Finsler structures in geometric mechanics and gravity, DGDS Monographs, Vol. 7 (Geometry Balkan Press, 2006) arXiv:gr-qc/0508023.
- [25] R. Coquereaux, “Clifford algebras, spinors and fundamental interactions: Twenty years after,” *Adv. Appl. Clifford Algebras* 19, 673-686 (2005), arXiv:math-ph/0509040.
- [26] C. Castro, “The extended relativity theory in Born-Clifford phase spaces with a lower and upper length scales and Clifford group geometric unification,” *Found. Phys.* 35, 971-1041 (2005).
- [27] A. Garrett Lisi, “Clifford bundle formulation of BF gravity generalized to the standard model,” (2005), arXiv:gr-qc/0511120.
- [28] M. Pavsic, “Spin gauge theory of gravity in Clifford space,” *J. Phys. Conf. Ser.* 33, 422-427 (2006), arXiv:gr-qc/0511124.
- [29] C. Castro, “The exceptional E₈ geometry of clifford (16) superspace and conformal gravity Yang-Mills grand unification,” *Int. J. Geom. Meth. Mod. Phys.* 6, 385-417 (2009).
- [30] F. Smarandache and V. Christianto, “Kaluza-Klein-Carmeli Metric from Quaternion-Clifford Space, Lorentz’ Force, and Some Observables,” (2008).
- [31] C. Castro, “The Clifford space geometry of conformal gravity and U(4) x U(4) Yang-

- Mills unification,” *Int. J. Mod. Phys. A* 25, 123-143 (2010).
- [32] M. Pavsic, “On the unification of interactions by Clifford algebra,” *Adv. Appl. Clifford Algebras* 20, 781-801 (2010).
- [33] C. Castro, “A Clifford $Cl(5, C)$ Unified Gauge Field Theory of Conformal Gravity, Maxwell and $U(4)$ Yang-Mills in 4D,” *Adv. Appl. Clifford Algebras* 22, 1-21 (2012).
- [34] C. Castro, “Lanczos-Lovelock and $f(R)$ gravity from Clifford space geometry,” *Int. J. Theor. Phys.* 52, 28-41 (2013).
- [35] C. Castro, “Lanczos-Lovelock-Cartan gravity from Clifford space geometry,” *Int. J. Geom. Meth. Mod. Phys.* 10, 1350019 (2013).
- [36] W. A. Rodrigues and S. A. Wainer, “A Clifford Bundle Approach to the Differential Geometry of Branes,” *Adv. Appl. Clifford Algebras* 24, 817-847 (2014), arXiv:1309.4007 [math-ph].
- [37] C. Castro, “A Clifford algebra-based grand unification program of gravity and the Standard Model: a review study,” *Can. J. Phys.* 92, 1501-1527 (2014).
- [38] C. Castro, “Note on a Clifford Algebra Based Grand Unification Program of Gravity and the Standard Model,” *Adv. Appl. Clifford Algebras* 26, 573-575 (2016).
- [39] C. HWANG and H. LEE, “Fermi Fields, Clifford Algebras and Path Integrals,” *New Phys. Sae Mulli* 66, 742-747 (2016).
- [40] C. Castro, “Clifford Algebraic Unification of Conformal Gravity with an Extended Standard Model,” *Adv. Appl. Clifford Algebras* 27, 1031-1042 (2017).
- [41] O. C. Stoica, “The Standard Model Algebra - Leptons, Quarks, and Gauge from the Complex Clifford Algebra Cl_6 ,” *Adv. Appl. Clifford Algebras* 28, 52 (2018), arXiv:1702.04336 [hep-th].
- [42] O. C. Stoica, “Leptons, Quarks, and Gauge Symmetries, from a Clifford Algebra,” *Springer Proc. Math. Stat.* 255, 309-316 (2017).
- [43] M. Pavšič, “Quantized Fields ‘a la Clifford and Unification,” (2017), arXiv:1707.05695 [physics.gen-ph].
- [44] W. Lu, “A Clifford algebra approach to chiral symmetry breaking and fermion mass hierarchies,” *Int. J. Mod. Phys. A* 32, 1750159 (2017), arXiv:1709.09294 [hep-ph].
- [45] M. A. Kurkov and F. Lizzi, “Clifford Structures in Noncommutative Geometry and the Extended Scalar Sector,” *Phys. Rev. D* 97, 085024 (2018), arXiv:1801.00260 [hep-th].
- [46] Y.-Q. Gu, “Clifford Algebra, Lorentz Transformation and Unified Field Theory,” *Adv. Appl. Clifford Algebras* 28, 37 (2018).
- [47] N. Gresnigt, “Sedenions, the Clifford algebra $Cl(8)$, and three fermion generations,” *PoS EPS-HEP2019*, 615 (2020).
- [48] N. G. Gresnigt, “The Standard Model particle content with complete gauge symmetries from the minimal ideals of two Clifford algebras,” *Eur. Phys. J. C* 80, 583 (2020), arXiv:2003.08814 [physics.gen-ph].
- [49] D. Lukman, M. Komendyak, and N. M. Borstnik, “Relations between Clifford algebra and Dirac matrices in the presence of families,” *Particles* 3, 518-531 (2020), arXiv:2004.00421 [physics.gen-ph].
- [50] N. G. Gresnigt, “A topological model of composite preons from the minimal ideals of two Clifford algebras,” *Phys. Lett. B* 808, 135687 (2020), arXiv:2004.11140 [physics.gen-ph].
- [51] M. A. Trindade, S. Floquet, and J. D. M. Vianna, “Clifford algebras, algebraic spinors, quantum information and applications,” *Mod. Phys. Lett. A* 35, 2050239 (2020), arXiv:2005.04231 [quant-ph].
- [52] I. Sogami and K. Koizumi, “Renovation of the Standard Model with Clifford-Dirac algebras for chiral triplets,” *Phys. Lett. B* 808, 135622 (2020).
- [53] W. Drechsler, “ $U(4)$ spin gauge theory over a Riemann-Cartan space-time,” *Z. Phys. C* 41, 197-205 (1988).
- [54] H. Dehnen and E. Hitzer, “Spin gauge theory of gravity with Higgs field mechanism,”

- Int. J. Theor. Phys. 33, 575-592 (1994), arXiv:1306.2085 [gr-qc].
- [55] Z.-H. Weng, “Four interactions in the sedenion curved spaces,” Int. J. Geom. Meth. Mod. Phys. 16, 1950019 (2018), arXiv:1812.02535 [physics.gen-ph].
- [56] F. W. Hehl, P. von der Heyde, G. D. Kerlick, and J. M. Nester, “General relativity with spin and torsion: Foundations and prospects,” Reviews of Modern Physics 48, 393-416 (1976).
- [57] M. Tecchiolli, “On the Mathematics of Coframe Formalism and Einstein-Cartan Theory—A Brief Review,” Universe 5, 206 (2019), arXiv:2008.08314 [math-ph].
- [58] O. Cebecioglu and S. Kibaroglu, “Maxwell-Modified Metric Affine Gravity,” (2021), arXiv:2104.12670 [hep-th].
- [59] J. C. Pati and A. Salam, “Lepton Number as the Fourth Color,” Phys.Rev. D10, 275-289 (1974).
- [60] J. Baez, “An Introduction to spin foam models of quantum gravity and BF theory,” Lect. Notes Phys. 543,25-93 (2000), arXiv:gr-qc/9905087.
- [61] C. Isham and J. Butterfield, “Some possible roles for topos theory in quantum theory and quantum gravity,” Found. Phys. 30, 1707-1735 (2000), arXiv:gr-qc/9910005.
- [62] K. A. Frøyshov, “Equivariant aspects of Yang-Mills Floer theory,” Topology 41, 525-552 (2002).
- [63] I. P. Zois, “Noncommutative topological quantum field theory-Noncommutative Floer Homology,” (2005), arXiv:hep-th/0510005.
- [64] A. Doring and C. Isham, “A Topos foundation for theories of physics. I. Formal languages for physics,” J. Math. Phys. 49, 053515 (2008), arXiv:quant-ph/0703060.
- [65] A. Doring and C. Isham, “A Topos foundation for theories of physics. II. Daseinisation and the liberation of quantum theory,” J. Math. Phys. 49, 053516 (2008), arXiv:quant-ph/0703062.
- [66] A. Doring and C. Isham, “A Topos foundation for theories of physics. III. The Representation of physical quantities with arrows,” J. Math. Phys. 49, 053517 (2008), arXiv:quant-ph/0703064.
- [67] K. Fukaya, “Floer homology of Lagrangian submanifolds,” (2011), arXiv:1106.4882 [math.SG].
- [68] M. Bochicchio, “Yang-Mills mass gap, Floer homology, glueball spectrum, and conformal window in large-N QCD,” (2013), arXiv:1312.1350 [hep-th].
- [69] C.-H. Cho and H. Hong, “Finite group actions on Lagrangian Floer theory,” J. Sympl. Geom. 15, 307-420 (2017), arXiv:1307.4573 [math.SG].
- [70] I. P. Zois, “Noncommutative Topological Quantum Field Theory, Noncommutative Floer Homology, Noncommutative Hodge Theory,” in 2nd International Conference on Mathematical Modeling in Physical Sciences (2014) arXiv:1401.4080 [math-ph].
- [71] K. Nakayama, “Topos quantum theory on quantization-induced sheaves,” J. Math. Phys. 55, 102103 (2014), arXiv:1404.2370 [math-ph].
- [72] B. Jubin and P. Schapira, “Sheaves and D-Modules on Lorentzian Manifolds,” Letters in Mathematical Physics 106, 607-648 (2016), arXiv:1510.01499 [math.AG].
- [73] M. Benini, A. Schenkel, and U. Schreiber, “The Stack of Yang-Mills Fields on Lorentzian Manifolds,” Communications in Mathematical Physics (2018),10.1007/s00220-018-3120-1, arXiv:1704.01378 [math-ph].
- [74] C. Viterbo, “Sheaf Quantization of Lagrangians and Floer cohomology,” arXiv:1901.09440 (2019), arXiv:1901.09440.
- [75] T. Kuwagaki, “Sheaf quantization from exact WKB analysis,” arXiv:2006.14872 (2020).
- [76] T. Asano and Y. Ike, “Sheaf quantization and intersection of rational Lagrangian immersions,” arXiv:2005.05088 (2020).
- [77] S. Chern, W. Chen, and K. Lam, Lectures

- on differential geometry (1999).
- [78] M. Fecko, Differential geometry and Lie groups for physicists (2006).
- [79] D.-S. Li, “A fermions representation in Pati-Salam model,” (2020), arXiv:2011.14362 [hep-ph].
- [80] K. Olive et al. (Particle Data Group), “Review of Particle Physics,” Chin.Phys. C38, 090001 (2014), arXiv:1412.1408 [astro-ph.CO].
- [81] Y. Fukuda et al. (Super-Kamiokande Collaboration), “Measurements of the solar neutrino flux from Super-Kamiokande’s first 300 days,” Phys.Rev.Lett. 81, 1158-1162 (1998), arXiv:hep-ex/9805021[hep-ex].
- [82] P. Minkowski, “ $\mu \rightarrow e\gamma$ at a Rate of One Out of 109 Muon Decays?” Phys. Lett. B 67, 421-428 (1977).
- [83] R. N. Mohapatra and G. Senjanovic, “Neutrino Mass and Spontaneous Parity Nonconservation,” Phys. Rev. Lett. 44, 912 (1980).
- [84] T. Yanagida, “Horizontal Symmetry and Masses of Neutrinos,” Prog. Theor. Phys. 64, 1103 (1980).
- [85] J. Schechter and J. Valle, “Neutrino Masses in $SU(2) \times U(1)$ Theories,” Phys. Rev. D 22, 2227 (1980).
- [86] Z.-z. Xing and S. Zhou, Neutrinos in particle physics, astronomy and cosmology (2011).
- [87] D.-S. Li, “Minimal unified field theory from square root metric, fiber bundle theory and Sheaf theory,” arXiv:1412.2578 [hep-th], (2014).
- [88] H. Nikolic, “Violation of unitarity by Hawking radiation does not violate energy-momentum conservation,” JCAP 04, 002 (2015), arXiv:1502.04324 [hep-th].
- [89] J. Preskill, “Do black holes destroy information?” in International Symposium on Black holes, Mem-branes, Wormholes and Superstrings (1992) pp. 22-39, arXiv: hep-th/9209058.
- [90] J. Barbon, “Black holes, information and holography,” J. Phys. Conf. Ser. 171, 012009 (2009).
- [91] S. Hawking, “Particle Creation by Black Holes,” Commun. Math. Phys. 43, 199-220 (1975), [Erratum: Commun.Math.Phys. 46, 206 (1976)].
- [92] Z.-z. Xing, H. Zhang, and S. Zhou, “Updated Values of Running Quark and Lepton Masses,” Phys. Rev. D 77, 113016 (2008), arXiv:0712.1419 [hep-ph].
- [93] S. Weinberg, The Quantum theory of fields. Vol. 1: Foundations (1995).
- [94] S. Weinberg, “The quantum theory of fields. Vol. 2: Modern applications,” (1996).
- [95] J. Cornwell, Group theory in physics. Vol. 2 (1985).